



VACCINES	FALL 2020 Immunizations for School Attendance
DTaP/DT Tdap/Td Diphtheria, Tetanus, Pertussis	<p>Kindergarten Four (4) or more doses of DTaP or DT, or any combination. Three doses of Td or a combination of Td and Tdap is the minimum acceptable for children ages 7 and older, with proper spacing.</p> <p>Grades 1-12 Four (4) or more doses of DTaP or DT, or any combination. If all four doses were given before the fourth birthday, a fifth (5th) dose is required. If the fourth dose was administered at least six months after the third dose, and on or after the fourth birthday, a fifth dose is not required. *</p> <p>Grades 7-12 One (1) dose of Tdap vaccine must be administered on or after the 10th birthday. **</p>
POLIO	<p>K-10 Three (3) or more doses of IPV. The FINAL dose must be administered on or after the fourth birthday regardless of the number of previous doses. If a combination of OPV and IPV was received, four (4) doses of either vaccine are required. ***</p> <p>Grades 11-12 Three (3) or more doses of IPV or OPV. If the third dose of either series was received prior to the fourth birthday, a fourth (4th) dose is required; if a combination of OPV and IPV was received, four (4) doses of either vaccine are required.</p>
MMR Measles, Mumps, Rubella	<p>K-12 Two (2) doses of MMR. Dose one (1) must be administered on or after the first birthday. The second (2nd) dose must be administered at least 28 days after dose one (1).</p>
HEP B Hepatitis B	<p>K-12 Three (3) or (4) doses of hepatitis B. The second dose must be administered at least 28 days after the first dose. The third dose must be given at least 16 weeks after the first dose and at least eight weeks after the second dose. The last dose in the series (third or fourth dose) must not be administered before age 24 weeks.</p>
VARICELLA (Chickenpox)	<p>K-10 Two (2) doses of varicella vaccine must be administered prior to entry. Dose one (1) must be administered on or after the first birthday. The second dose should be administered at least three months after dose one (1); however, if the second dose is administered at least 28 days after the first dose, it is considered valid.</p> <p>Grades 11-12 One (1) dose of varicella vaccine must be administered on or after the first birthday.</p>
MCV4 Meningococcal	<p>Grades 7-11 One (1) dose of meningococcal (serogroup A, C, W, and Y) vaccine must be administered prior to entry.</p> <p>Grade 12 Two (2) doses of meningococcal (serogroup A, C, W, and Y) vaccine must be administered prior to entry. ****</p>

NOTES:

- Vaccine should be administered according to the most recent version of the *Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger* or the *Catch-up Immunization Schedule for Persons Aged 4 Months Through 18 Years Who Start Late or Who Are More Than 1 Month Behind*, as published by the Advisory Committee on Immunization Practices. Schedules are available for print or download at <https://www.cdc.gov/vaccines/schedules/index.html>.
- Vaccine doses administered ≤ 4 days before the minimum interval or age are valid (grace period). Doses administered ≥ 5 days earlier than the minimum interval or age are not valid doses and should be repeated as age-appropriate. If MMR and varicella are not given on the same day, the doses must be separated by at least 28 days with no grace period.
- For additional information, please refer to the Ohio Revised Code 3313.67 and 3313.671 for School Attendance and the [ODH Director's Journal Entry](#) (available at <http://www.odh.ohio.gov>, [Immunization: Required Vaccines for Child Care & School](#)). These documents list required and recommended immunizations and indicate exemptions to immunizations.
- **Please contact the Ohio Department of Health Immunization Program at (800) 282-0546 or (614) 466-4643 with questions or concerns.**

* Recommended DTaP or DT minimum intervals for kindergarten students four (4) weeks between doses 1-2 and 2-3; six (6) month minimum intervals between doses 3-4 and 4-5. If a fifth dose is administered prior to the fourth birthday, a sixth dose is recommended but not required.

** Tdap can be given regardless of the interval since the last tetanus or diphtheria toxoid-containing vaccine. Children age 7 years or older with an incomplete history of DTaP should be given Tdap as the first dose in the catch-up series. If given at age 7-9 years, the routine Tdap dose at age 11-12 years should be given. If given at age 10 years, no additional dose is needed at age 11-12 years.

*** The final polio dose in the IPV series must be administered at age 4 or older with at least six months between the final and previous dose.

**** Recommended MCV4 minimum interval of at least eight (8) weeks between dose one (1) and dose two (2). If the first (1st) dose of MCV4 was administered on or after the 16th birthday, a second (2nd) dose is not required. If a pupil is in 12th grade and is 15 years of age or younger, only one (1) dose is required. Currently, there are no school entry requirements for meningococcal B vaccine.